

DEUTERIUM ARRAY MEMO #055

**MASSACHUSETTS INSTITUTE OF TECHNOLOGY**  
**HAYSTACK OBSERVATORY**  
 WESTFORD, MASSACHUSETTS 01886  
 September 20, 2004

*Telephone: 978-692-4764*  
*Fax: 781-981-0590*

To: Deuterium Array Group

From: Alan E.E. Rogers

Subject: The theoretical D line strength

The expected D1 line strength is estimated from the quantum mechanics of the D1 atom. In practice we are only interested in the relative opacities of D1 and H for a given abundance ratio  $n_D/n_H$  however, lets work from first principles. The probability of spontaneous decay, also frequently called the "Einstein A" coefficient can be calculated from (see Condon and Shortley equation 7<sup>4</sup> (3))

$$A = \frac{64\pi^4 \nu^3 M |\mu_0|^2}{3hc^3 g_a}$$

where  $\nu$  = frequency Hz  
 $h$  = Plank's constant =  $6.626 \times 10^{-27}$  erg s  
 $c$  = velocity of light =  $2.9979 \times 10^{10}$  cm s<sup>-1</sup>  
 $\mu_0^2 = \left( \frac{eh}{4\pi mc} \right)^2 = 8.6 \times 10^{-41}$  erg<sup>2</sup> gauss<sup>-2</sup>  
 $g_a$  = upper state degeneracy =  $2F_a + 1$

M is the sum of matrix elements squared.

For hydrogen:

S = 1/2 electron spin  
 I = 1/2 proton spin

And the vector F = I + S takes on the value of 0 from the ground state and 1 for the upper 3 states whose projections  $M_F = -1, 0, 1$  are 21 cm above the ground state.

The matrix elements squared (from Townes equations 5-65 and 5-66) are

$$|\mu|^2 = \frac{4 \left[ (I + 1/2)^2 \right] \mu_0^2}{(2I + 1)^2} \text{ when } \Delta m_F = 0$$

and

$$|\mu|^2 = \frac{2(I + \frac{1}{2})(I + \frac{3}{2})\mu_0^2}{(2I + 1)^2}$$

and all 3 possible transitions have values of  $|\mu|^2 = |\mu_0|^2$  so that

$A = 2.86 \times 10^{-15}$  for Hydrogen

And this value can be found in many places and to my knowledge has never been in dispute.

For deuterium

$$\begin{aligned} S &= \frac{1}{2} \text{ electron spin} \\ I &= 1 \text{ deuteron spin} \end{aligned}$$

For deuterium F takes on the values of  $\frac{1}{2}$  for the ground states and  $\frac{3}{2}$  for those states 327 MHz above ground. The ground state splits into a doublet with  $m_F = -\frac{1}{2}, \frac{1}{2}$  and the upper state momentum vector takes on 4 projections  $m_F = -\frac{3}{2}, -\frac{1}{2}, \frac{1}{2}, \frac{3}{2}$  and the matrix elements are no longer just equal to the Bohr magneton  $\mu_0$  squared.

The values are given by

$$\frac{4 \left[ \left( \frac{3}{2} \right)^2 - m_F^2 \right]}{9} \text{ for } \Delta m_F = 0$$

$$\frac{2 \left( \frac{3}{2} \pm m_F \right) \left( \frac{5}{2} \pm m_F \right)}{9} \text{ from } \Delta m_F = \pm 1$$

and take on the values of

$$\begin{aligned} 8/9 &\text{ for } \Delta m_F = 0, |m_F| = \frac{1}{2} \\ 4/3 &\text{ for } |\Delta m_F| = 1; m_F = \frac{1}{2} \\ 4/9 &\text{ for } |\Delta m_F| = 1; m_F = -\frac{1}{2} \\ g_a &= 2(3/2) + 1 = 4 \end{aligned}$$

for a sum over all 6 possible transitions of  $16/3$

so that

$$A_D = \frac{64\pi^4 \nu^3 \mu_0^2 (16/3)}{12hc^3} = 4.6 \times 10^{-17} \text{ sec}^{-1}$$

This value has been in contention.

Shklovsky (page 258) gives a value of  $6.6 \times 10^{-17} \text{ sec}^{-1}$  and a relative opacity ratio  $\tau_D/\tau_H \approx 0.4 \quad n_D/n_H$  while Weinreb points out errors in Shklovsky's equations and quotes a new value of  $4.65 \times 10^{-17} \text{ sec}^{-1}$  which comes from Field. The value assumed by Weinreb results an opacity ratio of  $\tau_D/\tau_H \approx 0.3 \quad n_D/n_H$ .

Sandy Weinreb also mentions that Field's calculation has been checked by Alan Barrett.

The optical depth  $\tau$  is given by

$$\tau = \frac{hc^2 A_{a \rightarrow b} g_a}{8\pi k \nu T_{ex} \sum g} \int_0^L N f(\nu, e) d\nu$$

where  $T_{ex}$  = excitation temperature

So that the ratio  $\tau_D/\tau_H$  is given by

$$\frac{\tau_D}{\tau_H} = \left( \frac{g}{\sum g} \right)_D \left( \frac{\sum g}{g} \right)_H \left( \frac{A_D}{A_H} \right) \left( \frac{\nu_H}{\nu_D} \right)^2 (n_D/n_H)$$

on the assumption that the line is only broadened by Doppler so that the Doppler widths are proportional to frequency.

With the revised value of  $A_D$  the ratio is reduced to  $\tau_D/\tau_H \approx 0.27 \quad (n_D/n_H)$ .

In a more recent paper Gould calculates the Einstein A coefficients for H and D and gets a value of  $A = 4.69675 \times 10^{-17} \text{ sec}^{-1}$  in his table 1, which is close the value of Field. Gould includes some small corrections to A which presumably result in a more accurate value. Using Gould's values the opacity ratio is  $(\tau_D/\tau_H) = 0.272(n_D/n_H)$ .

References:

Condon and Shortley, The theory of atomic spectra

Townes – Townes and Schawlow, Microwave Spectroscopy, Mc Graw- Hill, 1995

Shklovsky – I.S. Shklovsky, Cosmic Radio Waves, Harvard Univ. Press, 1960.

Field – Excitation of the Hydrogen 21 cm line, Proc. IRE, **46**, pp240-250, 1958.

Weinreb – S. Weinreb, A digital spectral analysis technique and its application to radio astronomy MIT Technical report #412, 1963.

Gould – R. J. Gould, Radiative Hyperfine Transitions, Ap.J. 423, pp. 522-528, 1994.